

## FACT SHEET

**'Fish Creek'** *Salix purpurea*

'Fish Creek' is a fast-growing shrub willow. This high-yielding variety is disease resistant, produces multiple small stems, and is well suited for biomass plantings, snowfences, streambank restoration, and riparian buffers.





**Botanical Name:** *Salix purpurea* 'Fish Creek'  
(Family: Salicaceae)

U.S. Plant Patent 17,710  
issued May 8, 2007

**Hardiness:** U.S.D.A. Zones 4 - 6

**Development:** 'Fish Creek' was produced through controlled willow breeding in 1998 as part of a research project to produce new willow cultivars that generate high biomass yields on a variety of sites, display resistance to diseases and pests, and possess agronomic traits suitable for mechanical planting, harvesting, and post-harvest processing. 'Fish Creek' was produced by crossing *Salix purpurea* '94006' with *S. purpurea* '94001'.

**Significance:** 'Fish Creek' is a shrub willow cultivar displaying exceptionally rapid growth, producing 4-5 dry tonnes per acre annually in yield trials, with low incidence of rust disease or damage by beetle or sawfly. Woody stems can be harvested every three to four years, and new shoots will re-sprout the following season. Repeated harvesting of shrub willow plantations can be sustained for at least 15 years.

**Description:**

**Height and Width:** 15-20 feet tall, 3-5 foot crown spread at 3 years when grown at 2 x 3 foot spacing.

**Habit:** Fast-growing, deciduous shrub with multiple small-diameter, vertical stems.

**Foliage:** Dark green oblong leaves, typically 2-3 inches long, 0.5-1 inches wide, with foliage April through October in Zone 5.

**Bark:** Brown when young, turning green and smooth with age; yellow-orange buds in winter.

**Flowers:** Male, early spring.

**Seeds:** No seeds produced.

**Culture:** Adaptable to a wide range of soil and moisture conditions. Prefers maximum sunlight.

**Propagation:** Roots easily from dormant stem cuttings.

**Uses:** Excellent for bioenergy plantations, streambank restoration, living snowfences, and riparian buffers. Dried stems can be used in basketry.

**Availability:** Available from Double A Willow ([www.doubleawillow.com](http://www.doubleawillow.com))

Originally published 2007 by SUNY-ESF

Kimberly D. Cameron<sup>1</sup>, Lawrence Smart<sup>1\*</sup>, Benjamin Ballard<sup>2</sup>, Timothy Volk<sup>3</sup>, and Lawrence Abrahamson<sup>3</sup>

<sup>1</sup>Dept. of Horticultural Sciences, <sup>2</sup>SUNY at Morrisville, NY 13408, <sup>3</sup>SUNY College of Environmental Science & Forestry, Syracuse, NY 13210

\*Contact: [lbs33@cornell.edu](mailto:lbs33@cornell.edu), 315-787-2490

Funding provided by New York Farm Viability Institute to LB Smart.

This variety is licensed by the research foundation of SUNY exclusively to Double A Willow

